MAKES WORK EASIER

EASY TO HANDLE
Soft-tack hyaluronan adhesion ensures easy application.
Does not require positioning during dressing changes.

CUT YOUR OWN SHAPE
Sorelex may be cut according to the shape of the wound.

LESS DRESSING CHANGES
According to the requirements and condition of the wound, the Sorelex primary dressing may be changed every two-five days. It is recommended to change the secondary dressing according to the exudate production

CONTIPRO

Sorelex

CLEANSES & DRIVES HEALING



OCTENIDINE
&
HYALURONIC ACID

CONTIPRO

Contipro a.s.
Czech Republic

+420 465 519 530
sales@contipro.cz

www.hyaluronanhealing.com

SORELEX **CLEANSES AND DRIVES HEALING**

Sorelex **effectively** establishes the optimal wound healing environment and **proactively** initiates wound healing processes, including the support of angiogenesis

- MANAGES INFECTION, DESLOUGHS, CLEANSES
- REDUCES HIGH LEVELS OF EXUDATE
- GENERATES NEW TISSUE GROWTH

SYNERGY OF **TWO ACTIVE SUBSTANCES**

The combination of a natural ingredient with unique regenerative properties and a safe and potent antiseptic ingredient provides rapid and visible results.

	WOUND BED PREPARATION	**TISSUE** REGENERATION
OCTENIDINE	Safely cleanses the wound Effective against G+ve/ G-ve bacteria No bacterial resistence	Protects from development of infection No inhibition of healing processes Low cytotoxicity to skin cells, no wound residues
HYALURONAN	Regulates the process of inflammation Initiates tissue regeneration process Stimulates healing processes	Supports viability and migration of keratinocytes and fibroblasts Improves vascularisation Regulates formation of collagen
VISIBLE EFFECT	Wound bed is promptly cleansed Exudation and malodour reduced Granulation tissue starts to be formed	New granulation tissue grows quickly Epithelialisation commences Scarring is reduced

FOR **COMPLICATED WOUNDS**

Sorelex provides solution to a wide spectrum of wounds. It is especially beneficial when rapid wound bed preparation and/or initiation of the healing processes are needed. **It is recommended for wounds with medium to high levels of exudate production**.

- INFECTED, SLOUGHY WOUNDS
- WOUNDS AT RISK OF INFECTION
- CHRONIC NON-HEALING WOUNDS

SOPHISTICATED CONCEPT



TRANSMISSION LAYER
Absorbs and transfers excessive exudate out of the wound into a secondary dressing. Prevents maceration of the skin around the wound. The layer has a distinctive red print; it should be applied AWAY FROM THE WOUND.

DESLOUGHING LAYER
Attracts wound slough, which is removed during dressing change.

ACTIVE LAYER
The unique combination of the active ingredients octenidine and hyaluronic acid having both antimicrobial and biological impact. A gel is formed in the wound. The active layer should be applied INTO THE WOUND.

SAFE HEALING

The active ingredients contained in the Sorelex dressing are **completely safe, with no adverse effects, even with long-term use**.
Thanks to its safety, Sorelex may be used even in risk populations (children, pregnant or breast-feeding women, diabetics or allergic patients).